

5040 N. 15th Avenue, Suite 307 . Phoenix AZ 85015 Tel 602 759 1245 . Fax 602 759 8331 . www.CyrexLabs.com

ACCESSION #: 13-113250 REQUISITION #: T10131437 SAMPLE TYPE: Serum **DATE COLLECTED:** 10/24/2013 **DATE RECEIVED:** 10/24/2013 **DATE OF REPORT:** 10/30/2013

PATIENT

Name: QC, JESSICA 5-3

DOB: 01/01/1900

Gender: M

PRACTITIONER

DEMODR, DEMODR TEST

5040 N. 15th Avenue Suite 307

Phoenix, Arizona 85015

TEST		RESULT			
Array 5 – Multiple Autoimmune Reactivity Screen **	IN RANGE (Normal)	EQUIVOCAL*	OUT OF RANGE	REFERENCE (ELISA Index)	
Parietal Cell + ATPase	0.73			0.1-1.4	
Intrinsic Factor		1.01		0.1-1.2	
ASCA + ANCA			2.07	0.2-1.4	
Tropomyosin ****		1.38		0.1-1.5	
Thyroglobulin	0.80			0.1-1.3	
Thyroid Peroxidase	0.94			0.1-1.3	
21-Hydroxylase (Adrenal Cortex)			1.25	0.2-1.2	
Myocardial Peptide		1.23		0.1-1.5	
Alpha-Myosin	1.11			0.3-1.5	
Phospholipid			2.04	0.2-1.3	
Platelet Glycoprotein		1.19		0.1-1.3	
Ovary/Testis ***		1.00		0.1-1.2	
Fibulin			1.60	0.4-1.6	
Collagen Complex	1.14			0.2-1.6	
Arthritic Peptide		1.21		0.2-1.3	
Osteocyte			1.73	0.1-1.4	
Cytochrome P450 (Hepatocyte)	1.21			0.3-1.6	
Insulin + Islet Cell			2.04	0.4-1.7	
Glutamic Acid Decarboxylase 65		1.25		0.2-1.6	
Myelin Basic Protein		1.15		0.1-1.4	
Asialoganglioside			1.44	0.1-1.4	
Alpha-Tubulin + Beta-Tubulin	0.61			0.4-1.4	
Cerebellar		1.13		0.2-1.4	
Synapsin		0.96		0.1-1.2	

^{****} Please note that the reference range for Tropomyosin ONLY has been updated with effect from 10/01/2013.

Dhiren Joshipura, M.D., Medical Director

Cyrex Laboratories is certified under the Clinical Laboratory Improvement Amendments of 1988 ("CLIA") as qualified to perform high-complexity clinical testing. Test result data on its own does not constitute a diagnosis. Only a physician or qualified healthcare professional should interpret the significance of a clinical lab test or make a diagnosis. This test was developed and its performance characteristics determined by Cyrex Laboratories, LLC. The names and titles of tests and arrays are for reference purposes only.

^{***} Ovary and Testis are tested together to avoid any confusion arising out of potential cross-reactivity

^{**} All analytes are tested for IgG and IgA combined.

^{*} Reference ranges are calculated based on the mean ±2 standard deviations (SD). Results > 1 SD, and <2 SDs above the mean are considered to be equivocal. An equivocal result represents the range between negative and suspicious low positive results. Results >2 SDs are considered out of range, and positive.